

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Radiation Toxins – Effects of Radiation Toxicity, Molecular Mechanisms of Action, Radiomimetic Properties and Possible Countermeasures for Radiation Injury

Dmitri Popov<sup>1</sup>, Jeffrey Jones<sup>2</sup> and Vacheslav Maliev<sup>3</sup>

<sup>1</sup>*Advanced Medical Technologies & Systems,  
Richmond Hill, Ontario, L4E4W8,*

<sup>2</sup>*Baylor College of Medicine, One Baylor Plaza, Houston, Texas,*

<sup>3</sup>*Vladicaucasian Scientific Center of Russian Academy of Sciences,  
Biotechnology Dept., Kosta Hetagyrova, Vladicaucas, North Ossetia–Alania,*

<sup>1</sup>*Canada*

<sup>2</sup>*USA*

<sup>3</sup>*Russia*

## 1. Introduction

### Radiation toxins induce high toxicity reactions after exposure

Acute Radiation Disease (ARD) or Acute Radiation Syndromes (ARS) are defined as the collective toxic clinical states observed from the acute pathological processes in various doses of irradiated mammals; to include: systemic inflammatory response syndrome (SIRS), toxic multiple organ injury (TMOI), toxic multiple organ dysfunction syndromes (TMODS), and finally, toxic multiple organ failure (TMOF). [2, 10, 18, 21 ] Moderate and high doses of radiation induces necrosis of radiosensitive cells with the subsequent formation of radiation toxins and their induced acute inflammatory processes. Radiation necrosis is the most substantial and most severe form of radiation induced injury, and when widespread, has grave therapeutic implications [1, 3, 53]. Low doses of radiation exposure induces apoptosis (controlled, programmed death of radiosensitive cells) without significant levels of specific radiation-induced toxin formation and with only low levels of inflammatory response [17, 50, 62].

Studying the trigger mechanism for radiation-induced lymphocyte death, N.I. Sorokina used the results of numerous experiments to show that ionizing radiation induces changes in the antigen phenotype of immature thymocytes in mice. This has the same type of effect as chemical differentiation inducers and thymotropin, which indirectly attests to the specific modifying effect of ionizing radiation [97, 98].

B.D. Zhivotovskiy described radiation-induced apoptosis and demonstrated a quantitative association between the pyknotic changes in the cell nuclei of thymocytes and production of postradiation chromatin decay products. The enzyme responsible for the decomposition of chromatin in irradiated cells is Ca/Mg-dependent endonuclease. Areas of endonuclease

attack are distributed randomly and are not associated with the level of repetition of nucleotide sequences of DNA or the transcriptional activity of chromatin. This supports the conclusion that the radiation death of lymphoid cells and the general biological phenomenon of programmed cell death are identical in nature [95, 96]. Apoptosis, interphase cell death of irradiated lymphocytes, is considered one example of the biological phenomenon known as "programmed cell death". Interphase cell death occurs as the result of "switching on" a specific genetic program, the triggering of which evokes induction of specific genes that switch on the "cell-death program". [1] Apoptosis may occur as the result of many signals, among which include: various types of lympholytic agents, different chemical agents, and physical factors which include ionizing radiation [15,50].

But relatively high doses of radiation could induce other forms of programmed cell death, apoptosis and/or necrosis. Necrosis especially, initiates inflammation and formation of specific Radiation Toxins (RTs) [62, 64, 65, 66, 89, 90, 91]. Specific Radiation Toxins are playing an important role as the trigger mechanism for inflammation, cell lysis, and damage to vital cellular structures such as mitochondria, DNA, ion channels and cell membranes [89, 90, 91, 92, 93, 94]. The mechanism and action of Radiation Toxins could be compared to actions of different microbial toxins, snake and scorpion venoms, enteric bacterial toxins, natural and plant-borne toxins [6, 86, 87, 88]. At high doses of radiation, specific Radiation Toxins play an important role in the development of pathological processes, especially injury to the central nervous system [89, 90, 91]. Radiation induces changes in concentration of many chemical mediators which possess significant influence on the Central Neural System – acetylcholine, cholinesterase, adrenergic amines, glutamate,  $\gamma$ -Aminobutyric acid, aspartic acid and interact with receptors and ions channels.[4, 30]. Many natural toxins like scorpion venom induce a prolongation of action potentials caused by selective inhibition of sodium current inactivation and specifically affect voltage-gated sodium channels in excitable tissues.[6, 29, 36, 44, 49, 60, 82, 83, 84, 85]. "Excitotoxicity" is an important pathological mechanism which damages the central nervous system [27,105]. After high doses of radiation, some specific receptors such as the NMDA receptor and AMP receptor are over activated. "Excitotoxins" such as NMDA and kainic acid, which bind to these receptors, as well as pathologically high levels of glutamate, can cause excitotoxicity, by allowing high levels of calcium ions to enter the cell.  $\text{Ca}^{++}$  influx into cells activates a number of enzymes, including phospholipases, endonucleases, and proteases such as calpain. These enzymes go on to damage cell structures e.g. components of the cytoskeleton, membranes, and DNA. Excitotoxicity may be involved in stroke, traumatic brain injury and neurodegenerative diseases of the central nervous system (CNS) such as multiple sclerosis, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), fibromyalgia, Parkinson's disease, and Huntington's disease.[27, 105]. It therefore seems logical that excitotoxicity possibly plays a prominent role in CNS injury.

We also postulate that specific Radiation Neurotoxins damage cerebral blood vessels and can induce activation of specific neurotransmitters which may produce exitotoxicity and its consequences [100, 103, 104]. The general toxicity of different types of ionizing radiation is associated mainly with a formation of Specific Radiation Toxins - a group of Radiation Toxins (RT) – Specific Radiation Determinants (SRD). [100, 103, 104]. Radiation toxins concentrate initially in intercellular fluid then disperse into blood and lymph and thereby induce development of systemic injury of hematologic and lymphatic systems. The toxicity of RT SRD subsequently distributes to the cerebrovascular, cardiovascular, and gastrointestinal systems, depending on the magnitude of the exposure. The injury to

cellular components of radiosensitive and pluripotent hematopoietic and other stem cells are induced by processes that are specific for different form of ARS. [100, 103, 104 ].

*Radiation Toxins:* The specific Radiation Toxins (group SRD) are composed from glycoproteins with high enzymatic activity including high proteolytic, lipidolytic, carbohydratelytic properties. RT are formed and accumulated in mammalian cells and migrate into blood and lymphatic circulation in the first hours after irradiation. Radiation Toxins, for the most part, are glycoproteins, pro-enzymes which are activated after relatively high doses of irradiation, i.e. after some radiation threshold for activation is exceeded. This group of glycoproteins has been isolated from cells of irradiated mammalian organisms, and possess the ability to induce inflammatory reactions, apoptosis, and necrosis similar to reactions mediated by radiation itself. The molecular weight of RT (SRD-group) range from 200-250 kDa. The SRD molecules have been isolated from blood, lymph and cells of irradiated animals but lymph has the highest concentrations measured. Irradiated animals develop the different forms of Acute Radiation Syndromes such as Cerebrovascular ARS, Cardiovascular ARS, Gastrointestinal ARS, depending on the level of radiation they were exposed to.

Acute Radiation Syndromes	Type of Radiation Toxins	Grade ARS Mild	Grade ARS Moderate	Grade ARS Severe	Grade ARS Extremely Severe
Cerebral form of ARS	cerebro-vascular R.T.	SRD-1.1	SRD-1.2	SRD-1.3	SRD-1.4
Cardio-vascular form of ARS	cardio-vascular R.T.	SRD-2.1	SRD-2.2	SRD-2.3	SRD-2.4
Gastro-intestinal form of ARS	gastro-intestinal R.T.	SRD-3.1	SRD-3.2	SRD-3.3	SRD-3.4
Hematological form of ARS	hemato-poietic R.T.	SRD-4.1	SRD-4.2	SRD-4.3	SRD-4.4

Table 1. Classification of radiation toxins (SRD group): Radiation Toxins are composed of toxic substances isolated from lymph of irradiated animals with different forms of Acute Radiation Syndromes.

*RT Properties-* In our experiments RT were isolated from central lymphatics of irradiated animals with different form of ARS and each of the RT possess differences in their radiomimetic properties. Cerebrovascular RT, and to a lesser extent, cardiovascular RT and gastrointestinal RT are neurotoxins. Hematopoetic RT – are strong hematotoxins. Hematopoetic RT can destroy red blood cells and initiate hemolysis, disrupt the blood clotting system and cause multi-organ degeneration and tissues damages. Hematopoietic RT possess important activity against pluripotent stem cells and blood marrow.

Component (%)	Toxic substance SRD-1	Toxic substance SRD -2	Toxic substance SRD-3	Toxic substance SRD-4.
Protein	44.9 ± 0.05	48.2 ± 0.03	50.1± 0.09	56.2 ± 0.12
Lipid	40.1± 0.04	39.6 ± 0.05	38.2± 0.04	30.1± 0.09
Carbohydrate	13.4 ± 0.09	11.3 ± 0.07	10.2 ± 0,03	10.1 ± 0,07
Mineral residue	1.6 ± 0.07	0.9 ± 0.02	1.3 ± 0.04	3.4± 0.17

Table 2. Biochemical composition of the toxic substances comprising Radiation Toxins. (SRD- Specific Radiation Determinants)

2. Material and methods

These studies were conducted at several different research institutions and laboratories listed as follows: Biotechnology Centre of Russian Academy of Science (North Osetia), Kazan All-Union Scientific Research Veterinary, Institute Belarussian Scientific and Research Institute for Radiobiology in Gomel, the St. Petersburg Veterinary Institute, the Advanced Medical Technology and Systems Inc., Ottawa-Richmond Hill, Ontario, Canada. The studies were approved by the Animal Care and Use Committee for ethical animal research equivalent, at each institution. A largest volume of purified Radiation Toxin was prepared from larger mammalian irradiated animals. Subsequently the RT were characterized chemically and biologically. The experimental design of later studies compared relative toxicity, potential for development of acute radiation syndromes, and potential lethality after intravenous or intramuscular injections of SRD containing Radiation Toxins. More recent studies focused on the immunogenicity of small doses of SRDs and the potential to use these agents as vaccine countermeasures against radiation injury.

2.1 Experimental animals

These experiments have employed a wide variety of experimental and agricultural animals which include : Black motley cattle, 2.5-3.0 years of age, live weight of 300-350 kg (n=134); Ukrainian pigs, 0.5-1.0 years of age, live weight 35-90 kg (n=142); “prekos” sheep, 3-12 months old, live weight 18-23 kg (n=156); mixed breed (mongrel) dogs, 2-4 years of age, live weight 6.0-6,5 kg (n=162), Chinchilla rabbits, 11-12 months old, live weight 3.5-3.7 (n=180), Latvian draft horses, 3-8 years of age, live weight 350-550 kg (n=32), Balb mice, 2-3 months old, live weight 20-22 g (n=2636), Wistar rats, 3-4 months old, live weight 180-220 g (n=4002). In all animals, the pre-experiment blood profiles, body temperature, central venous and lymphatic pressure did not exceed the limits of normal variability, (which had been previously measured for five species of mammals. Throughout the entire period of each experiment, ranging from 1 day to 2 years, the control and experimental animals were maintained under identical conditions of feeding, housing, and care, corresponding to livestock maintenance requirements and standards.



## 2.2 Electromagnetic radiation exposure

The animals were irradiated in RUM-17, Puma, and Panorama devices. The dose rate varied from 30Gy to 100Gy

## 2.3 Experimental design

Four independent experiments were performed with administration (IV or IM) of Radiation Toxins (SRD RT) to healthy, radiation naive animals that induced development of clinical symptoms of the Acute Radiation Syndromes. The administration of injection of the toxin was administered into veins (IV), into a muscle (IM), into the abdomen (IP) or into the skin (SC).

Experiment N1. Administration of Cerebrovascular form of Radiation Toxins to radiation naive animals in doses 0.1 mg/kg; 0,5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg up to 30 mg/kg initiates development of specific toxic reactions with symptoms mimicking the cerebral form of ARS. Injection of Cerebrovascular Radiation Toxins (SRD-1) in excess of 3 mg/kg doses to rats, rabbits, sheep resulted in lethality.

Experiment N2. Administration of radiation toxins of Cardiovascular Radiation Toxins (SRD-2) to radiation naive animals in doses 0.1 mg/kg; 0,5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg up to 30 mg/kg produced specific toxic reactions with symptoms mimicking the Cardiovascular form of ARS.

Experiment N3. Administration of Gastrointestinal Radiation Toxins (SRD-3) group to radiation naive animals in doses 0.1 mg/kg; 0,5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg up to 30 mg/kg produced development of specific toxic reactions with symptoms mimicking the GI form of ARS.

Experiment N4. Administration of Haemotopoietic Radiation Toxins (SRD-4) to radiation naive animals in doses 0.1 mg/kg; 0,5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg up to 30 mg/kg produced specific toxic reactions with symptoms of the hematological form of ARS.

## 2.4 Radiation toxins description and isolation

The methods of immune depletion, affine immuno-lympho-plasmasorption, as well as direct extraction were used to refine and purify the specific Radiation Toxins from the central lymph of animals with Cerebrovascular, Cardiovascular, Gastrointestinal and Haemotopoietic forms of Radiation Toxins.[104, 110]

Specific Radiation Determinants (SRD)- a group of Radiation Toxins isolated from lymph of irradiated mammals, had been divided to four important groups: 1.Cerebrovascular neurotoxic RT (SRD-1); 2.Cardiovascular RT(SRD-2); 3.Gastrointestinal RT (SRD-3); 4.Hematopoietic RT (SRD-4). The radiomimetic properties of Radiation Toxins preparations were evaluated on the basis of their capacity to induce radiobiological effects in radiation-naive animals after they were administered parenterally. Specific Cerebrovascular, Cardiovascular, Gastrointestinal and Hematopoietic Toxins (isolated from the lymph of animals irradiated at doses inducing Cerebral, Cardiovascular, Gastrointestinal and Haemotopoietic clinical forms of ARS) were dissolved in an isotonic solution of NaCl. The doses of preparations of Radiation Toxins which were administered was based on computation of the amount of SRD per unit volume of central lymph and absorbed dose of radiation. Radiation toxins of SRD group possess highly toxic properties and induce neuro-vascular and hematotoxic reactions in short time after administration.

Neurotoxins of SRD-1 group isolated from L.S. (lymphatic system) of irradiated animals with Cerebrovascular ARS after intra-muscular or intra-venous single injection in doses 5 mg/kg, 10 mg/kg, 15 mg / kg, 30 mg/kg activate acute neuro-vascular reactions with signs of injury of central and peripheral nervous system and damage of vascular walls of central nervous system.

Neurotoxins of SRD-2 group isolated from L.S. of irradiated animals with cardio-vascular form of ARS after intra-muscular or intra-venous single injection in doses 5 mg/kg, 10 mg/kg, 15 mg/kg, 30 mg/kg activate acute cardio-vascular and neuro-vascular reactions with signs of injury of central and peripheral vascular system. Neurotoxins of SRD-3 group isolated from L.S. of irradiated animals with gastro-intestinal form of ARS after intra-muscular or intra-venous single injection in doses 5 mg/kg, 10 mg/kg, 15 mg/kg and 30 mg/kg induce acute gastrointestinal syndrome with signs of injury of vascular system of gastro-intestinal system and injury of epithelium of intestinal walls. Hematotoxin of SRD-4 isolated from L.S. of irradiated animals with Hematopoietic form of ARS after single dose injection with doses 5 mg/kg, 10 mg/kg, 15 mg/kg, 30 mg/kg activate hematotoxic reactions with activation red blood cells and white blood cell lysis and activation of apoptosis of hematopoietic cells progenitors.

### 3. Results

#### **SRD-1 Radiation Toxins induced injury of Central Nervous System and development of Acute Cerebrovascular form of Radiation Syndrome**

The Cerebrovascular form of Radiation Toxins (SRD-1) were administered to radiation naïve animals in doses of 0.1 mg/kg; 0.5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg and 30 mg/kg. The injection of Cerebrovascular Radiation Toxins (SRD-1) to rats, rabbits, sheep produced bioeffects similar to the cerebrovascular form of ARS. (see Table 3) After parenteral administration, the animals experienced a short period of extreme agitation, followed by a deep coma, with alterations in breathing patterns and subsequently cardiovascular collapse. The results of autopsy of their bodies demonstrated cerebral hemorrhagic strokes, cerebrospinal fluid with blood (reddish color), hemorrhagic lesions in brain tissue. Internal organs were filled with blood. Multiple petechiae were observed on serous membranes. Depending on doses of cerebrovascular Radiation Neurotoxins, death was registered in 15 min or up to 5 hours after injection.

Case #1: The species studied in the experiment- 4 month old lambs. Radiation neurotoxin (SRD-1) in single dose of 5 mg/kg of weight was administered intramuscularly to radiation-naïve sheep. The injected animals were dead within 20 minutes after injection, with a profound clinical picture of cerebrovascular ARS. Within 5 minutes there was central nervous excitation and nervousness, and within 15 minutes there were signs of coma, followed by death. Post-mortem histopathology revealed: Brain- red discoloration of cerebrospinal fluid (CSF) with inflammatory meninges; multiple petechiae and hemorrhage on meninges surfaces. Grey matter of the brain - hemorrhages. Liver- dark-red coloration, with multiple petechiae and hemorrhages. Kidneys- multiple petechiae and hemorrhages. Clinical Diagnosis: Acute Cerebrovascular Shock / Acute Cerebrovascular form of ARS induced by single injection of radiation neurotoxin

Radiation Toxins	Symptoms	Toxic doses 5mg/kg	Toxic doses 10mg/kg	Toxic doses 15mg/kg	Toxic doses 30 mg/kg
Cerebrovascular Neurotoxin SRD-1.	Cerebral edema, hemorrhagic stroke	Death in one –three hours	Death in one hour	Death in 15-30 min	Death in 15 min
Cardiovascular SRD-2.	Cardiac insufficiency, hemorrhagic stroke.	Death in three days	Death in 14 hours	Death in 7 hours	Death in 2 hours
Gastro-Intestinal SRD-3.	Vomiting, diarrhea, hematochezia and melana.	Death in 10-15 days	Death in 5-7 days	Death in 2-3 days	Death in 24 hours
Hematopoietic SRD-4	Peripheral edema, ecchymosis, anemia, granulocytopenic fevers.	Death in 2-4 weeks.	Death in next 10 days.	Death in next 7-10 days.	Death in next 4-5 days.
Mixture of lower doses of toxins SRD 1+2+3+4	Cerebral edema, vomiting, hemorrhage, cardiac insufficiency.	Death in 4-5 hours	Death in 1-3 hours	Death in one hour	Death in one hour

Table 3. Radiation Toxins: Toxic properties of Radiation Toxins (SRD Group 1-4 ) after single injection and different forms of Acute Radiation Syndromes induced by each.

**SRD-2 Radiation Toxins induced injury of the Cardiovascular System and development of symptoms mimicking the Acute Cardiovascular Radiation Syndrome**

The Cardiovascular form of Radiation Toxins were administered to radiation naive animals in doses of 0.1 mg/kg; 0,5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg and 30 mg/kg. The injection of Cardiovascular (SRD-2) RT in toxic doses to rats, rabbits, sheep produced symptoms similar to the cardiovascular form of ARS. (see Table 3) In this experiment, the following symptoms were observed: a delayed, shorter duration and less extreme period of agitation than with SRD-1 RTs, followed by tachycardia, cardiac arrhythmias, tachypnea. Postmortem histopathological analysis revealed inflammatory changes in the cardiac muscle tissue.

Case #2: Species in experiment- 4 months old lamb. Radiation neurotoxin (SRD-2) in single dose 5 mg/kg of weight was administered intravenously. 14 hours after injection all the injected lambs were dead after with a clinical picture of cardiovascular shock: agitation, fever, agitation, tachycardia, arrhythmias, hypotension, hyperventilation, followed by lethargy, signs of coma and death. Post-mortal histopathology showed: Heart- damaged microvasculature, cardiac vasoconstriction , ischemic myocardium, contraction band necrosis, multiple cardiac hemorrhages. Kidneys: multiple petechies of haemorrhages, acute tubular necrosis and renal failure.



### **SRD-3 Radiation Toxins induced injury of the Gastrointestinal System and development of symptoms mimicking the Acute Gastrointestinal Radiation Syndrome**

Administration of Gastrointestinal Radiation Toxins (SRD-3) were administered to radiation naive animals in doses of 0.1 mg/kg; 0.5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg and 30 mg/kg. Injection of SRD-3 in toxic doses to experimental animals produced different grades of the Gastrointestinal form of ARS. (see Table 3) Case #3: Experimental species: one year old sheep. After a single dose injection of 15 mg/kg of the SRD-3 Gastrointestinal radiation toxins (isolated from central lymph of animals with the gastrointestinal form of ARS) intravenously, the animals elicit the following: Clinical signs-decrease in appetite and increased peristalsis within hours after injection, followed by severe diarrhea, hematochezia, dehydration, and later followed by collapse over several days. Death occurred 5-10 days after toxin administration. Histopathology revealed- local vasoconstriction, damaged microvasculature, large vessel vasodilatation and hyperemia of the small bowel and colon. Hypertension was manifested approximately two hours after intravenous injection. Ultimately hypotension developed after days of vomiting, diarrhea and dehydration.

### **SRD-4 Radiation Toxins induced development of Acute Hematopoietic Acute Radiation Syndrome**

The Hematopoietic Radiation Toxins (SRD-4) were administered to radiation naive animals in doses of 0.1 mg/kg; 0.5 mg/kg; 1 mg/kg; 2 mg/kg; 3 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg and 30 mg/kg. Injection of SRD-4 to experimental animals resulted in erythrocytopenia, lymphocytopenia, leukocytopenia, and thrombocytopenia within days to weeks after injection. (see Table 3) The development of clinical features of the Acute Hematopoietic Syndrome depended on the dose of SRD-4 Hematopoietic Radiation Toxins injected to radiation naïve animals. Autopsy of those animals that died showed acute or chronic hematotoxic reactions. The clinical signs were: short-term agitation within 2 hours after administration accompanied by a short-term leukocytosis which gave way to a progressive, profound leukopenia, mainly attributable to a decrease in the absolute number of lymphocytes, the minimal levels of which were measured between days 7 and 15 after injection. Total WBC was 1.2-1.6 1000/ $\mu$ l to 0.4-0.5 thousand/ $\mu$ l in sheep and 1.8-2.5 thousand/ $\mu$ l to 0.5-0.7 thousand/ $\mu$ l in cattle. The recovery of absolute and relative levels of leukocytes and lymphocytes was observed in some animals between days 30 and 60. Blood counts exhibited thrombocytopenia accompanied by progressive erythrocytopenia, which developed into profound anemia. An extensive blood analysis of the peripheral blood of the cattle showed that the processes induced by the SRD-4 injection and the processes occurring after irradiation were nearly identical. Analysis of the clinical reaction to SRD-4 administration, which was assessed on the basis of body temperature, and heart and respiration rate, established that all experimental animals showed reactions of the same type for all the tested doses and that sheep and horses were more sensitive to the administered preparation. Thus, in sheep, which received SRD-4 at the maximum dose tested, the body temperature increased by 1.5-2 ° C, reaching 41.2° 1 or 2 days before death. This core body temperature increase was accompanied by severe tachycardia and tachypnea, which were measured at 105-106/min. and 70-80/ minute, respectively. When SRD-4 was administered in the intermediate and maximum dose levels, both sheep and cattle showed changes of the same type but of lesser degree depending on the dose. The majority of experimental animals recovered between days 30-60 after injection of the preparation. Post-mortem examination of the animals that died showed changes characteristic of acute radiation

sickness, accompanied by marked hemorrhage. Death often will occur secondary to overwhelming bacterial or fungal sepsis. Some sheep showed areas of skin epilation on the back and abdomen.

Case# 4. Species- sheep. Hematoxin, SDR-4, isolated from irradiated mammals with the Hematopoietic form of ARS, was injected via single dose to non-irradiated mammals. The SRD-4 RT induced significant changes in white blood cells (WBC) and red cells profile (RBC). Hematoxin injected to non-irradiated sheep in doses 5 mg/kg, 10 mg/kg, 30 mg/kg activated a complex reaction which included general inflammation, vascular endothelial cell injury, apoptosis and necrosis of blood progenitor/ stem cells. The levels of sheep erythrocytes, leukocytes, lymphocytes, thrombocytes significantly increased in first hours and day after single injection of radiation hematotoxin and could be a result of general inflammation reactions and stimulation of immune system. However the levels of erythrocytes, leukocytes, lymphocytes, thrombocytes significantly decreased after 72 hours after hematoxin administration and the minimal level of erythrocytes, leukocytes, lymphocytes were measured after 168- 360 hours following a single dose injection of the radiation-induced hematoxin. (see Tables 4- 7) Characteristics of the peripheral blood of large horn cattle after a single injection of Hematopoietic Radiation Toxin (SRD-4) were measured and levels of erythrocytes, leukocytes, lymphocytes significantly increased in first hours after a single dose injection of radiation hematotoxin and decreased after 72, 168, and 360 hours after administration. (see Tables 8- 11)




Fig. 1. Dose-dependent changes in sheep peripheral blood erythrocytes after a single injection of Hematopoietic Radiation Toxin (SRD-4).




Fig. 2. Dose-dependent changes sheep peripheral blood leukocytes after a single injection of Hematopoietic Radiation Toxin (SRD-4).




Fig. 3. Dose-dependent dynamics of sheep Lymphocytes after single IV administration of Hematotoxin (SRD-4).




Fig. 4. Dose-dependent dynamics of sheep thrombocytes (platelets) after single IV administration of Hematotoxin (SRD-4).




Fig. 5. Dose-dependent characteristics of large horn cattle peripheral blood erythrocytes after single injection of Hematopoietic Radiation Toxin (SRD-4).




Fig. 6. Dose-dependent changes in cattle peripheral blood leukocytes after a single injection of Hematopoietic Radiation Toxin (SRD-4).




Fig. 7. Dose-dependent dynamics of cattle Lymphocytes after single IV administration of Hematotoxin (SRD-4).






Fig. 8. Dose-dependent dynamics of cattle thrombocytes (platelets) after single IV administration of Hematotoxin (SRD-4).

4. Discussion

Radiation toxin SRD-1 (cerebrovascular radio-neurotoxin) activates pathophysiological processes and clinical symptoms / signs of central and peripheral nervous system injury with development of neurological and cognitive deficit, activation, followed by depression, of specific centers of the central nervous system, including autonomic nervous system pathways.

Radiation toxin SRD-2 (cardiovascular radiotoxin ) activates pathophysiological processes and clinical symptoms / signs of cardiac and vascular inflammation and injury with endothelium cell, microvascular vessel and cardiovascular system tissue showing apoptosis and necrosis. Severe hypotension and tachycardia were important clinical indicators of cardiovascular acute radiation syndromes with the subsequent development of cardiac myocyte necrosis.

Radiation toxin SRD-3 (gastrointestinal radiotoxin) initiates symptoms signs of gastrointestinal system injury including anorexia and vomiting, induced by the toxic effects and damage to the vascular and lymphatic vessel network of the gastrointestinal system, as well as apoptosis of intestinal lining epithelial cells, including crypt / villi necrosis. Severe diarrhea and melena / hematochezia were important clinical indicators of the mimicked gastrointestinal acute radiation syndrome.

Radiation toxin SRD-4 (hematotoxin) induces development of red cell lysis and apoptosis of white blood cell and red cell progenitors. The action of the SRD-4 hematotoxin is described

in Tables 4-7 for sheep and 8-11 for cattle. Thrombocytopenia, lymphocytopenia, granulocytopenia, ecchymosis, hemorrhage and coagulopathy were important clinical signs of the mimicked hematopoietic acute radiation syndrome.

Other clinical indicators of the Radiation Toxins (RT) included cutaneous system involvement: including: cutaneous edemas, blistering, desquamation, hair loss, ulcer and necrosis.

Severe Acute Radiation Exposure syndromes will induce Toxic Multiple Organ Failure (TMOF) and Toxic Multiple Organ Involvement (TMOI) – e.g. pneumonitis, renal failure, renal hypo-perfusion, acute tubular necrosis, hepatic failure, etc. as an acute consequences of radiation toxemia. Similarly, bacterial toxemia induced by bacterial superantigens and radiation-induced toxemia are very similar in their clinical manifestation and similar in their pathophysiological actions. In a related matter, the molecular structure of radiation antigens (toxins) and bacterial superantigens, as well as different snake venom toxins have similarities.

Thus, if a researcher administers specific Radiation Toxins of one of the SRD groups to radiation naive animals, toxicity will be observed along a dose response curve, from doses as small as 0.1 mg/kg or 0.5 mg/kg up to the highest tested dose of 30 mg/kg. The observed toxicity will mimic the ARS syndrome specific to the exposure of the animals from which the SRDs were acquired. Thus the exogenously administered SRD initiates development of specific toxic reactions with symptoms of a specific ARS, hence the name-specific radiation determinant (SRD 1-4). The SRD-group molecules possess both toxic and antigenic properties. The molecular Koch's postulates, which are commonly applied to bacterial toxin pathogenicity, could also be applied to the biological effect produced by Radiation Toxins, and those postulates would mostly be satisfied to list RT as the causative agents [101, 106].

#### **4.1 Comparative analysis of bacterial toxins vs radiation toxins (see table 12)**

We compared lethal toxicity and clinicopathological features and characteristics for Radiation Toxins and Bacterial Toxins. Pathogenic bacteria which colonize the intestine, can invade intestinal epithelial cells and lymphatics, and/or produce one or more toxins that are important etiologies of diarrheal disease [32, 42, 48, 57, 86, 87, 88]. Bacteria produce toxins and the mechanisms of action of bacterial toxins can be classified into three important groups (see Table 12) : 1. Toxins with intrinsic enzymatic activity – cytotoxic bacterial toxins. 2. Toxins that binds to receptors that stimulate the actions of second messengers – cytotoxic bacterial toxins (e.g. botulinum or cholera toxins) . 3. Toxins that damage eukaryotic cell membranes – membranotoxic bacterial toxins. [32, 42, 48, 57, 86,]. We postulate that radiation toxins possess high enzymatic activity, damage cell membranes, and also activate cell receptors. We postulate that radiation toxin mechanisms of action depend on the toxin type and dose of radiation toxins. We postulate that Radiation Toxins could initiate two different types of cell death – programmed (apoptosis) or necrosis. [90,91].

Escherichia Coli is an Enteric Gram-negative rods and their pathogenic strains contained the wide group of toxins with different properties. Escherichia Coli contain endotoxin - lipopolysaccharide, which is antigenic and virulence factor. [12, 48]. Escherichia Coli contains three groups of antigens - O, H and K. The O antigens - cell wall antigens - found in polysaccharide portion of LPS. The H antigens are associated with flagella and the K antigens are associated with the fimbriae or capsule. [12,48,78,79,87,88 ]. Enterotoxigenic E.Coli - EAEC, produce three important toxins. EAEC heat stable enterotoxin is a 4.1-kDa protein. The

other EAEC heat -liable enterotoxin with molecular weight 120 kDa. The third EAEC enterotoxin is a heat labile,108 kDa protein with the ability to induce a severe acute intestinal inflammation. [12,78,79,87]. Enterohemorrhagic Escherechia coli, EHEC contains a Shigella-like toxin. The Shiga toxin family contains Shiga-like toxin 1 (SLT-1) and Shiga-like toxin 2 (SLT-2). EHEC strains produce SLT -2 or combination SLT 1 and SLT 2 [12 ]. Shiga toxin is a cytotoxic for endothelial and epithelial cells including human colonic and ileal epithelial cells, and B-lymphocytes [12]. Enteropathogenic Echerechia coli induce an important and distinctive histopathologic lesion in the intestine with destruction of microvilli – the classic histopathologic lesion - AE lesion [12 ]. There are two proteins responsible for development AE lesion – a 94-kDa protein and protein encoded by the eaeB gene. [12,78,79].

Toxins	Mechanism of Action	Symptoms
1. Enterotoxigenic E. Coli (ETEC).	1. ETEC - Three toxins: EA- ST, EA- HL, EA-HL	ETEC toxins induce hyper-secretion by intestinal mucosa cells, watery diarrhea
2. Enteropathogenic E. Coli (EPEC).	2. EPEC - Shiga-like toxins.	EPEC toxins causing destruction of microvilli and development focal lesions.
3. Enterohemorrhagic E. Coli (EHEC).	3. EHEC – verotoxin or Shiga-like toxin.	EHEC induce severe form of hemorrhagic colitis, acute renal failure or hemolytic uremic syndrome.
4. Enteroinvasive E. Coli (EIEC).		EIEC induce dysentery like syndrome
5. Enteroadherent E. Coli (EAEC).		EAEC cause traveler’ diarrhea

Table 4. Bacterial Toxins and their mechanisms of action.

Enterotoxigenic Escherechia coli produce heat – stable (ST) and heat-labile enterotoxins (LT). STa is a cysteine-rich, 18-amino-acid peptide. ST is produced as a precursor that is cleaved by signal peptidase and translocated to the periplasm where is added three intra-molecular disulfide bonds crucial for toxin activity. A second proteolytic agent occurs extracellularly and produce biologically active and toxic ST [12,81.]. STa acts by binding to a protein intestinal epithelial receptor localized in brush border membrane. [12,78]. E.coli heat-stable enterotoxin b is a 71-amino-acid precursor protein secreted extracellularly and proteolytically processed to a toxic form as a 48-amino-acid protein [12, 79]. E. coli heat-labile enterotoxin I is closely related to CT, including its structure and enzymatic activity. [12]. Other Escherechia coli Toxins include Cytolethal distending toxin and Cytotoxic necrotizing factors [12, 78,79 ]. Staphylococcus aureus produces serologically distinct protein exotoxins named A, B, C1, C3, D, E. All of these exotoxins are small proteins (24-30 kDa) with important biological activities. All seven toxins stimulate local receptors in upper intestinal tracts [12, 86]. Staphylococcal enterotoxins are likely playing an important role in cytokine production and are most potent T-cell receptor activators and inducers of lymphocyte proliferation. This class of toxins could be classified as “superantigens”, as well as the toxins produced by Streptococcus spp. and Clostridium perfringens. Superantigens presented directly to the T-

Clinical symptoms induced by Bacterial Superantigene	Clinical symptoms induced by radiation toxins (SRD group) and by irradiation.
Central Nervous System involvement Peripheral nervous system Confusion, Irritability, Lethargy	CNS involvement: Cerebro-Vascular Acute Radiation Syndrome. Radiation Induced Neurotoxin. Neurological and Cognitive Deficit, Vomiting. Severe form of brain microcirculation disorder. Clinical futures of hemorrhagic stroke.
Cardio-Vascular System involvement Systolic Blood Pressure :< 90 mmHg Tachycardia >100 beats/min, Hypotension, Hyperventilation, Loss of Sympathetic responsiveness .	Cardio-Vascular System involvement: Cardio-vascular Radiotoxin-induced Acute Radiation Syndrome.- Tachycardia, Bradicardia, Arrythmias, Hypertension followed by severe hypotension (Systolic Blood Pressure:< 80 mmHg), hyperventilation.
Gastro-Intestinal System Involvement Vomiting, diarrhea, hematochezia and stool mucous (depending on which enteric toxin).	Gastro-intestinal System Involvement: Gastro-Intestinal Acute Radiation Disease. Radiotoxin. Vomiting, diarrhea. Melena and hematochezia. Abdominal pain- moderate to severe.
Bacterial Superantigen hematotoxicity Thrombocytopenia, Platelet count < 100,000/mm3.- Thrombocytopenia., leukopenia, pancytopenia. Disseminated Intravascular Coagulation (DIC) Syndrome.	Hematopoietic Toxicity: Acute Radiation Hematopoietic Syndrome. Hematopoietic Radiation Toxin- Thrombocytopenia, Lymphocytopenia, Granulocytopenia. Pancytopenia.
Cutaneous involvement: diffuse maculopapular, petechial or ecchymotic rash, intense erythroderma, desquamation (e.g. meningococemia or toxic shock syndrome).	Cutaneous System Involvement: Possibleaction of any type of Radiation Toxins. Swelling, edema, blustering, desquamation, hair loss, ulceration and skin necrosis
Multiple Organ Failure and Multiple Organ Involvement.	Multiple Organ Failure and Multiple Organ Involvement.
Renal failure, Renal hypo-perfusion; Oliguria, Acute tubular necrosis	Renal failure, renal hypo-perfusion, Acute tubular necrosis.
Hepatic failure and inflammation	Hepatic failure
Pulmonary System: Hyperventilation with Respiratory alkalosis, pulmonary hypertension and edema, hypoxemia, ARDS	Pulmonary System: Hyperventilation. Radiation induced pneumonitis and subsequent pneumonia, ARDS.

Table 5. Comparative analysis of clinical symptoms induced by radiation and/or Radiation Toxins - (SRD group 1-4)- induced ARS and those produced by Toxic Shock Syndrome induced by Bacterial Superantigene.

cell receptors by cells containing major histocompatibility complex class II molecules, without processing by antigen-presenting cells [32,48,86]. Superantigens (SAGs) are a class of antigens which cause non-specific activation of T-cells resulting in polyclonal T cell activation and massive cytokine release. The large number of activated T-cells secrete large amounts of cytokines (the most important of which is TNF- $\alpha$ ). [32,86]. SAG stimulation of antigen presenting cells and T-cells elicits a response that is mainly inflammatory, focused on the action of TH-1 T-helper cells. Some of the major products are IL-1, IL-2, IL-6, TNF- $\alpha$ , gamma interferon (IFN- $\gamma$ ), macrophage inflammatory protein 1 $\alpha$  (MIP-1 $\alpha$ ), MIP-1 $\beta$ , and also monocyte chemoattractant protein 1 (MCP-1). Clinical symptoms / signs induced by bacterial Superantigens include central and peripheral nervous system involvement (confusion, lethargy, vomiting, anorexia), involvement of the cardiovascular system with development of hypotension, tachycardia and loss of sympathetic responsiveness. Bacterial Superantigens activate toxic processes within the gastrointestinal system including acute vomiting and diarrhea. The important signs and clinical picture of bacterial superantigen hemotoxicity with development thrombocytopenia, leukopenia, lymphocytopenia and development of coagulopathy. Bacterial Superantigens could activate injury of cutaneous system and diffuse rash, erythrodermia, desquamation. [32,86] Radiation Toxins of SRD groups 1-4 can activate similar pathological processes, clinical picture and clinical signs as the bacterial superantigen toxicity and the three entities: 1) Acute Radiation Syndromes, 2) SRD RT- induced acute syndromes and Bacterial Toxin- induced septic reaction and toxicity can be compared. (see Table 5)

#### **4.2 Comparative analysis of toxins produced by venom toxins and radiation toxins (see table 6)**

Neurotoxic proteins can be isolated from various snake venoms. These proteins can potentially be a mix of neurotoxins, which attack the nervous system; hemotoxins, which attack the circulatory system; plus cytotoxins, bungarotoxins and many other toxins that affect the body in different ways [6,29,49,60,61,82,83,84,85]. Almost all snake venom contains hyaluronidase, an enzyme that ensures rapid diffusion of the venom into the body of the bite-victim [82,84]. Viper venoms from the Elapidae, Hydrophiidae, Atractaspididae, Viperidae and Colubridae families contain at least 25 separate classes of biologically active compounds, including enzymes and non-enzymatic molecules. Venom- neurotoxins activate different reactions with nerve synapse – presynaptic and postsynaptic action; and induce anticholinesterase-induced flaccid paralysis. Myotoxins from Venom could induce systemic skeletal muscle damage. Hemotoxins induce the development of severe microcirculatory injury, damage to vascular membranes, thrombosis or bleeding. [6, 82, 84].

#### **4.3 Radiation induced apoptosis and necrosis**

Radiation induced Apoptosis and Necrosis as response of eukaryotic cells to influence of different types of radiation remain controversial. Radiation induced cell death by triggering apoptosis pathways was described in many articles and supported by many scientists [11, 15, 17, 39, 51, 52, 53, 54, 55, 56; 57, 58, 59, 62, 63, 74, 75, 76, 78]. However some scientists and some institutions described processes developing in irradiated eukaryotic cells as necrosis [24;34;38] and some scientists describing a variety of complex mechanism and mentioned that radiation-induced cell death could develop under different mechanisms of pathogenesis which include and apoptosis and necrosis. [1, 34, 38, 43, 77] Activation of



Venom Toxins	Mechanism of Action	Symptoms
Polypeptide toxins.	Autonomic, Neurotoxic, Cardiotoxic, Myotoxic	Induce two forms of programmed cell death: cell apoptosis or cell necrosis. Presynaptic or postsynaptic actions. Induce excitation of some neurons. Neurotoxicity, Haemototoxicity.
Phospholipase toxins.	Presynaptic neurotoxic, myolytic, procoagulant, cardiotoxic, necrotic.	Induce cerebrovascular toxicity, cardiovascular toxicity.
Enzyme toxins: Transferases, Oxydoreductases, Hydrolases, Carbon- nitrogen lyases, Metalloproteinases.	Disorder of haemostasis, necrotic, haemolytic.	Blood vessel injury, blood cell lysis, brain tissue damage, heart muscle damage, liver damage, kidney damage. Neurotoxicity. Haemototoxicity. Skin damage. Hair follicle damage.
The Group of Toxins: Lectins, Nerve growth factors, Phospholipase inhibitors, Complement inhibitors, Proteinase inhibitors.	Cardiotoxic, Neurotoxic.	Anticoagulant proteins. Platelet aggregation inducers. Platelet aggregation inhibitors.

Table 6. Venom toxins and mechanisms of action – biochemical and clinical classification.

cysteine proteases, cleavage of PARP - poly (ADP-ribosyl) ation of nuclear proteins causes the inner mitochondrial membrane to become permeable for ions and other small molecules during both types of cell death- apoptosis and necrosis. Many mechanisms of cell necrosis and apoptosis are identical – for example, degradation of the Nuclear matrix is a Common Element during Radiation-Induced Apoptosis and Necrosis [1,3,11,16,24,34,38,43,53,59,78]. Data from different sources described that the differences between apoptosis and necrosis are much less numerous than previously studies suggested. [1,78]. Joseph Dynlacht and his research group of scientists studied the outcome of human promyelocytic leukemia (HL60) cells irradiated with 10 or 50 Gy of X-rays and determined the mode of leukemic cell death. Different doses of X-rays induced different modes of cell death – cells irradiated with 10 Gy died by necrosis; cells irradiated with 50Gy died predominantly by apoptosis. [16]. The Cell Death Nomenclature Committee recommends the use of the appropriate diagnosis: apoptotic necrosis. [34]. Comparison and description of a cell’s Necrosis and Apoptosis were provided by a group of scientists from different institutions and research groups [16,34]. Apoptosis is described as a normally active, programmed process of cell death devoid of inflammation and toxins. [16,34]. Necrosis possesses the characteristics of accidental or externally-induced cell death from the influence of different environmental factors and triggering development of inflammation and elaborating of toxins, release of pro-inflammatory active and toxic cellular contents into the intracellular and extracellular fluids and into the lymphatic systems and blood circulation [39,55,56,62]. Caspases are important molecular mechanisms and central components of the apoptosis process [39, 50, 51, 52, 54, 58, 62, 74, 75, 77, 78].

Apoptosis	Necrosis
Inflammation never present.	Inflammation always present.
Toxic substances never present.	Toxic substances always present.
Morphologically cells shrinks, become denser, condensation occur, original name – shrinkage necrosis. Mitochondria’s structure and functions are not affected.	Morphologically cells swelling, lysis. Mitochondria’s structure and functions are affected.
Karyorhexis – pyknotic nuclear fragments. DNA broken into segments.	Karyolysis, Pyknosis, Karyorhesis. Nuclear swelling, Chromatin granular, Chromatin flocculation, Types of radiation induced damage of DNA: Breaks of the strand, alteration to bases, destruction of sugar, cross-links and formation of dimmers
Caspase activation always present.	Caspase depended initiation of developing of necrosis possible. Caspase-8 initiation possible with apoptosis and necrosis.
Genetic control initiate apoptosis	No genetic control, environmentally-induced
The Budding phenomenon. Cell membrane not affected	The Blebbing phenomenon, increased membrane’s permeability. Cell membrane affected by external and internal factors.

Table 7. Apoptosis and Necrosis: Comparison of Morphological and Biochemical features.

However, apoptosis is the principal mechanisms by which cells are physiologically eliminated. During apoptotic death, cellular dissociation by caspases of cell compartments occur and cell compartments and molecules are packaged into apoptotic bodies to avoid immune activation. [24,34,78]. Necrosis was considered in past times as passive, uncontrolled and unorganized mechanism of cell death. At the present time necrosis is considered as an alternative form of programmed cell death whose activation induces important biological consequences including immune reactions and induced inflammation. [43, 51, 52, 67, 74, 75, 77].

4.4 Inflammation induced by radiation

Radiation induces an important and aggressive inflammatory response in the irradiated tissues. [19, 20, 28, 30, 64, 65]. Acute Radiation Disease (ARD) or Acute Radiation Syndromes (ARS) were defined as a toxic, poisonous exposure with development of the acute pathological processes in irradiated animals: systemic inflammatory response syndrome (SIRS), toxic multiple organ injury (TMOI), toxic multiple organ dysfunction syndromes (TMOD), toxic multiple organ failure (TMOF) [2, 5, 18, 19, 21, 89, 90, 91]. Radiation-induced lung injury, radiation pneumonitis (RP), is a potentially fatal side-effect of thoracic radiation therapy [3, 4, 11, 15]. Radiation-induced thyroiditis: can occur after the use of radiation treatment and can result in lifelong hypothyroidism [18]. Irradiation of the liver with doses of 2,450-2,920 rads caused liver injury in 14 patients of 65 patients treated with radiation therapy [3, 28]. The clinical picture of radiation-induced hepatitis includes hepatomegaly, ascites, pleural effusion and alteration in liver function. Radiation injury of the central nervous system (CNS) has devastating and fatal clinical consequences and

severely limits the dose that can be delivered in the radiotherapy of brain, head and neck, pulmonary and other tumors [19, 28, 30, 64, 65]. Hematopoietic system toxicity is a major limiting factor in the use of aggressive, combined modality therapy (chemotherapy + radiotherapy) in the treatment of malignant disease [14, 22, 40].

Permanent anemia and pancytopenia can be caused by the reduced capability for cellular proliferation due to the stem cells injury produced by radiation. [14, 69, 70, 72, 76]. Radiation induces an inflammatory response in the irradiated organs characterized by leukocyte infiltration and vascular permeability changes and vascular injury [90, 99]. Vascular injury is a key determinant of acute and chronic organ intoxication and dysfunction associated with irradiation of the different systems, including the gastrointestinal tract [9, 19, 20, 22,]. Radiotherapy is used in the treatment of pediatric brain tumors and is often associated with inflammation and behavior changing following irradiation-induced inflammation and injury in the developing brain [19, 20]. The consequences of irradiation microglial loss can be secondary to injury from pronounced inflammation [19, 20]. A prominent inflammatory response occurs following both ischemic and hemorrhagic forms of microcirculation disorders [19, 20].

#### **4.5 Roles of radiation toxins in developing of cellular structure radiation-induced injury**

Radiation Toxins contained a group of biologically-active compounds which induce complicated but specific mechanisms of action with tissues and cellular damage. Radiation Toxins include a group of glycoproteins with high enzymatic activity against proteins, lipids, carbohydrates. Radiation Toxins contain a group of peptides with selective affinity to number of receptors and specific interactions with receptors of mammalian organisms. Toxins isolated from the irradiated mammals inhibit or activate a group of an ion channels, acetylcholine receptors, nicotinic receptors, membranes, coagulant/anticoagulant pathways. Our experiments involved injections to radiation-naïve animals of toxic doses of Radiation Toxins (SDR- group) which damaged endothelial cells of blood and lymphatic vessels, and injury to blood microcirculation in the brain, lungs, gastrointestinal tract, cardiovascular system. Administration (IV or IM) of the Haematopoietic Radiation Toxins to radiation-naïve animals induced development of anemia with reduction of hemoglobin concentration, reduction of red blood cell count and reduction of white cell blood count. Hematopoietic Radiation Toxins induced the development of hemolytic anaemia with accelerated destruction of mature red cells outside the bone marrow and destruction of pluripotent stem cells inside of the bone marrow. Radiation-induced hemolytic anemia occurred with intravascular and extravascular hemolysis. As a result of stem cell injury erythropenia, leukopenia, thrombocytopenia developed within days after irradiation. However, the acute hemolytic extravascular and intravascular anemia with lysis of erythrocytes, developed after high doses of radiation.

#### **4.6 Development of radiation countermeasures**

There are many sources of oxidative stress in the lives of workers, whether they work in nuclear power facilities, on the front lines of international conflicts, or in the reaches of outer space. The exposure dose can vary substantially, but at minimum will accelerate the aging of their organ systems, and at worse could result in acute exposure syndromes that may be fatal. A common thread of the oxidative stress exposures is ROS-binding to critical cellular

organelles and molecules, which can result in cellular dysfunction, mutation of nucleic acids, or even apoptotic cell death [107]. Currently there are no proven countermeasures for these exposures, aside from a clinical agent, Amifostine, (Etyhol™), which is used to reduce mucositis and other side effects from radiation therapy dose in cancer patients.

The cytotoxic effects of different types of radiation may be the single most important clinicopathologic process by which oxidative damage is induced from reactive oxygen species and radiation toxicity induced by radiation toxins. Radiation toxins (SRDs) with high enzymatic activity and their ability to degraded a wide variety of extracellular proteins, lipids, carbohydrates and DNA molecules, induce damage of important intracellular compartments such as mitochondria, ion channels, DNA, as well as activating degradation of peptide bonds in important polypeptides in tissues and vascular endothelium. Yet the exact mechanism by which radiation toxins stimulate development of the ARS is poorly understood.

Countermeasures against radiation toxicity can be developed both in oral formulas [109] and parenteral agents, e.g. Manganese SuperOxide Dismutase (MnSOD)-liposomes [108, 109] which can reduce radiation exposure-induced biological effects. The oral formulas consist of a combination of antioxidants and chemopreventive, anti-progression agents, which can reduce the likelihood of mild toxicity in acute low dose radiation exposures and tumor induction in chronic low dose exposures. The combination of oral formulas with higher dose reactive oxygen species scavengers, like Mn-SOD can improve survival in animals exposed to acute high dose total body irradiation [109]. In addition to these measures, active immunization by low, non-toxic doses of radiation toxins, (SRDs) can also be employed to reduce radiation toxicity. [104, 110] To be effective, SRD immunization must occur no less than 30 days before irradiation, and can be effective up to three years or more. Active immunization by radiation toxins can significantly improve the survival rate (up to 60%) versus placebo-controlled irradiated animals. Our studies attempt to show the potential ability of specific antibodies to neutralize radiation toxins and thus substantially reduce the effects on radiation-induced neuro-, vascular, gastrointestinal, and hematopoietic toxicity. [104, 110] Antiradiation antibodies prevent the radiation-induced cytolysis of selected groups of cells that are sensitive to radiation. Anti-radiation antibodies derived from different phases of the Acute Radiation Syndrome can compete with and thus prevent cytolysis mediated by cytotoxic lymphocytes. The therapeutic benefit of neutralization of SRD radiation toxins could make hemopoetic stem cell transplantation more effective. Antiradiation vaccine and IgG antibodies have shown activity in animals against several different types of radiation include gamma, heavy ions, and neutron irradiation.[104, 110]

Preliminary and pilot studies in vitro, in animal models and recently in humans, are showing some promise for both efficacy and safety/tolerability. [108-110] With current unpredictability in the operation of nuclear power facilities, potential radio-biologic terrorist weapons, and human operations in extreme environments, it is important to develop countermeasures to radiation toxicity in order to protect the potentially exposed. The first step in developing these countermeasures is to fully understanding the mechanism of toxicity, as is attempted to be summarized in this chapter.

## 5. Acknowledgements

Carlos Montesinos, Kedar Prasad, Michael Epperly, Joel Greenberger.



## 6. References

- [1] Akadi Y, et al. Radiation-induced Apoptosis and Necrosis in Molt-4 Cells: A Study of Dose-effect Relationships and Their Modification. *International Journal of Radiation Biology*. 1993, V.64, No 1, pp. 47-56.
- [2] Azizova T.V. et al. Multi-organ Involvement and failure in selected accident cases with acute radiation syndrome observed at the Mayak Nuclear Facility. *The British Institute of Radiology*.
- [3] Kumar A., Leeds N., Gregory N. et al. Malignant Gliomas: MR Imaging Spectrum of Radiation Therapy- and Chemotherapy-induced Necrosis of the Brain after Treatment.
- [4] Ahmed, A. et al. Effect of Exposure to Electromagnetic Radiation from the Mobile Phone on Acetylcholinesterase Activity in the Hippocampus and Striatum of Young and Adult Male Rats, *Med. J. Cairo Univ.*, 2006, 74(2),(Suppl.), 129-135.
- [5] Berger M. and al. Medical management of radiation injuries: current approaches. *Journal of Occupational Medicine*. 2006. V.56. pp.162-172.
- [6] Tilmisanuy A., Osman A. Snake, Scorpion and Envenomation: A Review. *JKAU: Med. Sci.*, Vol.3, pp. 11- 18.
- [7] Briggs S., Brinsfield K. *Advanced Disaster Medical response Manual for Providers*. Harvard Medical International, Boston, Ma.2003,
- [8] Blakely W. Early biodosimetry response: recommendations for mass-casualty radiation accidents and terrorism. *Armed Forces Radiobiology Research Institute, Uniformed Services University, Bethesda*. pp. 1-26.
- [9] Wong C.,Van der Kogel A. Mechanisms of Radiation injury to the Central Nervous System: Indications for Neuroprotection. *Molecular Interventions*. 10. Volume 4, Issue 5, 273-284, 2004.
- [10] Casarez E. Basic principles of toxicology. *BIOC*, 2001.p.597.Centers For Disease Control and Prevention. *Acute Radiation Syndrome: A fact Sheet for Physicians*. June 29, 2005. [www.bt.cdc.gov/radiation](http://www.bt.cdc.gov/radiation)
- [11] Chow, B., Li Y. and Wong, C. Radiation-induced apoptosis in the adult central nervous system is p53-dependent. *Cell Death and Differentiation* 7, 712-720 (2000).
- [12] Sears C, Kaper J. Enteric Bacterial Toxins: Mechanisms of Action and Linkage to Intestinal Secretion. *Microbiological Reviews*, Mar. 1996, p.167-215.
- [13] Netto D., Chiacchio S. et al. Humoral response and neutralization capacity of sheep serum inoculated with natural and cobalt 60 - irradiated *Crotalus durissus terrificus* venom.LAURETTI, 1968. *Journal of Venomous Animals and Toxins*. Vol. 8, No.2, 2002.
- [14] Daniak N, Ricks R. The evolving role of haemotopoetic cell transplantation in radiation injury: potential and limitations. *The British Institute of Radiology*, 2005. Pp. 169-174.
- [15] Dewey W et al. Radiation induced apoptosis: relevance to radiotherapy. *International Journal Radiation Oncology Biology Physics*. 1995, No 33, pp. 781-796.
- [16] Dunlacht J., et al. Degradation of the Nuclear matrix Is a common Element during Radiation-Induced Apoptosis and Necrosis. *Radiation Research*, 1999, v.152. No 6, pp.590 - 603.
- [17] Rio D., Pardo V. Monoamine neurotoxins-induced apoptosis in lymphocytes by a common oxidative stress mechanism: involvement of hydrogen peroxide



- (H<sub>2</sub>O<sub>2</sub>) caspase-3, and nuclear factor kappa-B (NF- kappa B), P53, c-Jun transcription factors. *Biochem Pharmacology Journal*, 2002, 63, pp. 677-688.
- [18] Fliedner T. et al. Multi-organ involvement as a pathogenic principle of the radiation syndromes: a study involving 110 cases histories documented in SEARCH and classified as the bases of hematopoietic indicators of effect. *The British Institute of Radiology*, 2005, p.1-6.
- [19] Kalm M. Inflammation and behavior following irradiation-induced injury in the developing brain. Center for Brain Repair and Rehabilitation, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Sweden, 2009.
- [20] Kleining T, Vink. R. Suppression of inflammation in ischemic and hemorrhagic stroke: therapeutic options. *Current Opinion in neurology*. 2009, No 22, pp. 294-301.
- [21] Konchalovsky M. et al. Multiple Organ Involvement and failure: selected Russian radiation accident cases re-visited. *The British Institute of radiology*. 2005,pp.26-29.
- [22] Swarts S. et al. Radiation-induced cyto-toxicity, DNA damage and DNA repair: implications for cell survival theory. *Radiation Environmental Biophysics*. 1990, 29 (2), pp.93 – 102.
- [23] Rakesh D. et al. Activation of a CrmA-insensitive, p35-sensitive pathway in Ionizing Radiation-induced Apoptosis. *The American Society for Biochemistry and Molecular Biology*. 1997, v.272, No. 3, pp. 1965 – 1969.
- [24] Fink S., Cookson B. Apoptosis, Pyroptosis and Necrosis: Mechanic Description of Dead and Dying Eukaryotic Cells. *Infection and Immunity*. Apr.2005, p.1907 – 1916.
- [25] Schultheiss, T., Kun, L., Ang, K. and Stephens, L. Radiation response of the central nervous system. *Int. J. Radiat. Oncol. Biol. Phys.* 31, 1093-1112 (1995).
- [26] Tofilon, P.J. and Fike, J.R. The radioresponse of the central nervous system: A dynamic process. *Radiation Research*, 153, 357-370 (2000).
- [27] Temple M., O'Leary D., Faden A. The role of glutamate receptors in the pathophysiology of traumatic CNS injury. Chapter 4 in *Head Trauma: Basic, Preclinical, and Clinical Directions*. Miller LP and Hayes RL, editors. Co-edited by Newcomb JK. John Wiley and Sons, Inc. New York. 2001. Pages 87-113.
- [28] Trot K. Radiation Risks from Imaging of Intestinal and Abdominal Inflammation. *Scandinavian Journal of Gastroenterology*. 1994, vol. 29, No 203, pp. 43-47.
- [29] Topchieva S., Mehrabova M., Influence of electromagnetic radiation on venom of *Vipera Lebetina* Obtuse. *Radioprotection*, 2008, vol. 43, n. 5.
- [30] Van Der Kogel, A. Radiation-induced damage in the central nervous system: An interpretation of target cell responses. *Br. J. Cancer Suppl.* 7, 207-217 (1986).
- [31] Hungerford J. Committee on Natural Toxins and Food Allergens. General Referee Reports: *Journal of AOAC International*. Vol., 89, No.1, 2006.
- [32] Rudbach J., et al. Preparation and Properties of a National Reference Endotoxin. *Journal of Clinical Microbiology*. Jan. 1976. P.21-25.
- [33] Luvisetto S. et al. Toxicity of botulinum neurotoxins in central nervous system of mice. *J.Toxicon*, 2003, March, 41(4), pp.475 – 481.
- [34] Levin S. Toxicological highlights: Apoptosis, Necrosis, or Oncosis: What is your Diagnosis? A Report from the Cell Death Nomenclature Committee of the Society of Toxicologic Pathologists. *Toxicological sciences*. 1998, 41, 155 -156.

- [35] MacCann D. Radiation poisoning: Current concepts in the Acute Radiation Syndrome. *American Journal of Clinical Medicine*. V.3, No3. 2006, pp. 13-21.
- [36] Ramirez-Dominguez M, Olamendi T. et al. Cn11, the first example of a scorpion toxin that a true blocker of Na<sup>+</sup> currents in crayfish neurons. *The Journal of Experimental Biology*, 205, 869-876(2002).
- [37] Molla M, Panes J. Radiation-induced intestinal inflammation. *World Journal of Gastroenterology*, 2007, No 13, pp. 3043-3046.
- [38] Guido M. and Isabelle J. Apoptosis, Oncosis, and Necrosis . An Overview of Cell Death. *American Journal of Pathology*. 1995, v.146, No 1.
- [39] Ozsahin M. et al., CD4 and CD8 T-lymphocyte Apoptosis Can Predict Radiation-Induced Late Toxicity: A Prospective Study in 399 Patients. *Clinical Cancer Research*, 2005, 11(20), October 15, 2005.
- [40] Xue M. et al. Intracortical Hemorrhage Injury in Rats : Relationship Between Blood Fractions and Brain Cell Death. *Journal of American Heart Association*, 2000, No.31, pp.1721-1727.
- [41] Nieder C. et al. Experimental concepts for toxicity prevention and tissue restoration after central nervous system irradiation. *Radiation Oncology*. 2007, No 2, p.23.
- [42] Sagara-Ishijima N., Furuhashi K. Toxic characteristics of the Synthetic Lipid A Derivative DT-5461 in Rats and Monkeys. *Toxicological Sciences*, 1999 ,49, 324-331.
- [43] Nieder C., Andratschke N., Price R., Rivera, B., and Ang, K. Innovative prevention strategies for radiation necrosis of the central nervous system. *Anticancer Research*, 22,1017-1023 .
- [44] Gilles N., Blanchet C. et al. A Scorpion  $\alpha$ -Like Toxin That Is Active on Insects and Mammals Reveals an Unexpected Specificity and Distribution of Sodium Channel Subtypes in Rat Brain Neurons. *The Journal of Neuroscience*, October 15, 1999.
- [45] Vladimirov A. Biological membranes and non-programmed cell death. *Soros educational Journal*. V.6, No 9, 2000, p.2-9.
- [46] Wasalenko J, McVittie T et al. Medical Management of the Acute Radiation Syndrome: Recommendations of the Strategic National Stockpile Radiation Working Group. 2004. *Ann Int Med*. No. 140, pp.1037-1051.
- [47] Bonner W. Low-dose radiation: Thresholds, bystander effects, and adaptive responses. *Proceeding of the national Academy of Sciences of United States of America*. April 29, 2003, vol.10, no. 9, pp. 4973-4975.
- [48] Ueno Y . Toxicology of microbial toxins. *Pure & Appl. Chem.*, Vol.58. No.2, pp.339-350. 1986.
- [49] Yu-Cheng X, Song-Ping L. Inhibition of Sodium Channels In Rat Dorsal Root Ganglion Neurons by Hainantoxin- IV, a novel Spider Toxin. *Acta Biochimica et Biophysica Sinica*, 2003, 35(1).
- [50] Watters D. Molecular mechanisms of ionizing radiation-induced apoptosis. *Immunology and Cell Biology* (1999) 77, 263-271.
- [51] Haimovitz-Friedman A. Radiation-induced signal transduction and stress response. *Radiat. Res*. 1998; 150: S102-8.
- [52] Szumiel I. Monitoring and signaling of radiation-induced damage in mammalian cells. *Radiat. Res*. 1998; 150: S92-101.

- [53] Kim Y, M.D. Radiation-induced Necrosis Deteriorating Neurological Symptoms and Mimicking Progression of Brain Metastasis after Stereotactic-guided Radiotherapy. *Cancer Res Treat.* 2007 March; 39(1): 16–21.
- [54] Waterhouse, N.J. et al. (2003) Mitochondrial outer-membrane permeabilization in apoptosis. In: *Cell Proliferation and Apoptosis*. Hughes, D. and Mehmet, H., eds. 185–200. (Bios Scientific Publishers, Ltd. Oxford).
- [55] Verheij M. et al. Radiation-induced apoptosis. *Cell Tissue Res.* 2000, Jul; 301 (1): 133–42.
- [56] Rajesh P. et al. Apoptosis: molecular mechanisms and pathogenicity. *EXCLI Journal* 2009;8:155-181 ISSN 1611-2156
- [57] Robertson J. and Orrenius S. Molecular Mechanisms of Apoptosis Induced by Cytotoxic Chemicals. *Critical Reviews in Toxicology*, 30(5):609–627 (2000).
- [58] Riedl S. et al. Molecular mechanisms of caspase regulations during apoptosis. *Molecular Cell Biology.* 2004, v.5. p.897–907.
- [59] Higuchi M. et al. Regulation of reactive oxygen species-induced apoptosis and necrosis by caspase 3-like proteases. *Oncogene.* 26 November 1998, Volume 17, Number 21, Pages 2753–2760.
- [60] Tan P, Khan A., Brusic V. *Bioinformatics for Venom and Toxins Science*. Henry Stewart Publications. 1467–5463. Briefings in Bioinformatics. Vol. 4. No 1. 53–62. March 2003.
- [61] Ji-Sheng C.. *An Assessment of Toxins*. Research Institute of Chemical Defense. P.O. Box 104. Beijing, 102205, China.
- [62] Kathryn D. Held. *Radiation Induced Apoptosis*. Massachusetts General Hospital. Harvard medical School.
- [63] Wirth B. and Olander D. *Fundamentals of Radiation Damage and Defects Production*. US Berkeley, Nuclear Engineering. Ref: Sect 17.2, D.O. text.
- [64] Molla M., Panes J. Radiation induced intestinal inflammation. *World Journal of Gastroenterology.* 2007. Pp.3043–3046.
- [65] Molla M. et al. Influence of dose-rate on inflammatory damage and adhesion molecule expression after abdominal radiation in the rat. *International Journal of Radiation Oncology and Biological Physics.* 1999.45. pp.1011–1018.
- [66] Linard C., et al. Acute induction of inflammatory cytokine expression after gamma-irradiation in the rat: effect of an NF-kappa B inhibitor. *International Journal of Radiation Oncology and Biological Physics.* 2004.58. pp.427–434.
- [67] Ramanan S. et al. Role of PPARs in Radiation-Induced Brain Injury. *PPAR Research.* V.2010. article ID 234975. 12 pages.
- [68] Schiller T., Chen Y. Modeling radiation-induced lung injury risk with an ensemble of support vector machines. *Journal Neurocomputing* , Volume 73. Issue 10–12, June, 2010.
- [69] Baum S. et al. Radiation Induced Anemia in Rats Exposed repeatedly to mixed Gamma-Neutron radiation. *Radiation Research.* 1970. 41. pp.492–499.
- [70] Dritshilo A. Sherman D. Radiation and Chemical Injury in the Bone Marrow. *Environment Health Prospectives.* 1981, v.39, pp. 59–64.
- [71] Wiebe E., and Rodrigues G. Radiation Induced Lung Injury. Strategies for reducing damages while optimizing therapeutic doses. 2006. Parchurst. Publisher for Oncology Exchange. V.5. No 2. pp.29–31.


- [72] Wei-Chung L. et al. Protection against Radiation-Induced Bone Marrow and Intestinal Injuries by *Cordyceps sinensis*, a Chinese Herbal Medicine. Radiation Research, 2006. 166, 900–907.
- [73] Wharton T. et al. Radiation Hepatitis induced by abdominal irradiation with the cobalt 60 moving strip technique. Presented as the Fifty-fourth Annual Meeting of the American Radium Society. 1973. pp.53-80.
- [74] Elinger A., Thomson C. death by Design: Death by Design : Apoptosis, Necrosis and Autophagy. Current Opinion in Cell Biology. 2004, pp. 653-669.
- [75] Dispersyn G. et al. Dissociation of cardiomyocyte apoptosis and differentiation in infarct border zones. European Heart Journal. 2002. 23. pp.849-857.
- [76] Ohgushi M. et al. Molecular Pathway and Cell State Responsible for Dissociation-Induced Apoptosis in Human Pluripotent Stem Cells. Cell Stem Cell 7, 225–239, August 6, 2010 Elsevier Inc.
- [77] Claudio A. et al. Caspase-dependent initiation of apoptosis and necrosis by the Fas receptor in lymphoid cells: onset of necrosis is associated with delayed ceramide increase. Journal of Cell Science 115, 4671-4683 © 2002 The Company of Biologists Ltd.
- [78] Rastogi P. et al. Apoptosis: Molecular Mechanisms and Pathogenicity. EXCLI Journal 2009;8:155-181 ISSN 1611-2156. Received: June 29, 2009, accepted: July 27, 2009, published: August 03, 2009.
- [79] Barnett N, editor. Biosecurity Reference: CFR-Listed Agent and Toxin Summaries. pp.1-150. Prepared by Sandia National Laboratory, Albuquerque, New Mexico 87185 and Livermore, California 94550.
- [80] Simon J. Nuclear, biological, and chemical terrorism: Understanding the threat and designing responses. Int J Emergency Mental health 1999; 1: 81-89.
- [81] Franz D, DVM, PhD, Colonel, Veterinary Corps. United States Army. Defense against Toxin Weapons. U.S. Army Medical Research and Materiel Command, pp. 1-36.
- [82] Shivaji P. Snake venom neurotoxins: Pharmacological classification. Journal of toxicology. Toxin reviews. 2004, vol. 23, n°1, pp. 37-96 [60 page(s)].
- [83] Clemetson K, Morita T., Kini M. Scientific and standardization committee communications: classification and nomenclature of snake venom C-type lectins and related proteins. Posted on ISTH website, November 19, 2008.
- [84] Viljorn C., Botes B. Snake Venom Toxins. The Journal of Biological Chemistry. Vol. 249, No. 2, Issue of January 25, pp. 366-372, 1974.
- [85] Oren B., Ph.D. Venom Toxins as Ion Channel Research Tools. Modulator No.21 Fall 2006. 28-31. [www.alomone.com](http://www.alomone.com)
- [86] Alouf, J. et al. 1991. The family of mitogenic, shock-inducing and superantigenic toxins from staphylococci and streptococci, p. 367–414. In J. E. Alouf and J. H. Freer (ed.), Sourcebook of bacterial protein toxins. Academic Press Ltd., London.
- [87] Fiorentini, C., et al. 1995. Escherichia coli cytotoxic necrotizing factor 1: evidence for induction of actin assembly by constitutive activation of the p21 Rho GTPase. Infect. Immun. 63:3936–3944.
- [88] Gariepy, J., et al. 1987. Importance of disulfide bridges in the structure and activity of Escherichia coli enterotoxin ST1b. Proc. Natl. Acad. Sci. USA 84:8907–8911.



- [89] Popov D., Maliev S. Antiradiation Vaccine: Immunological neutralization of Radiation Toxins at Acute Radiation Syndromes. 38th COSPAR Scientific Assembly. Held 18-15 July 2010, in Bremen, Germany, p.3.
- [90] Popov D., Maliev S. Radiation Toxins: Molecular mechanisms and radiomimetic properties. 38th COSPAR Scientific Assembly. Held 18-15 July 2010, in Bremen, Germany, p.3.
- [91] Popov D., Maliev S. Countermeasure development : Specific Immunoprophylaxis and Immunotherapy of Combined Acute Radiation Syndromes. 38th COSPAR Scientific Assembly. Held 18-15 July 2010, in Bremen, Germany, p.3.
- [92] Jones, J. A., Riggs, P. K., Yang, T., Pedemonte, C. H., Clarke, M, Feedback, D., and Au, W. Ionizing radiation-induced bioeffects in space and strategies to reduce cellular injury and carcinogenesis ASEM 78(4) Section II suppl., pp. A67-78, 2007.
- [93] Koniarek J, Thomas J., Vazquez V. Detection of microlesions induced by heavy ions using liposomes filled with fluorescent dye. *Advances in Space Research* , Volume 34, Issue 6, 2004, Pages 1373-1377. *Space Life Sciences: Radiation Risk Assessment and Radiation Measurements in Low Earth Orbit*.
- [94] Koniarek J., Worgul B., Do heavy ions cause microlesions in cell membranes? *Advances in Space Research*, Volume 12, Issues 2-3, 1992, Pages 417-420.
- [95] Zhivotovsky B., Samali, A., Gahm, A., and Orrenius, S Caspases: their intracellular localization and translocation during apoptosis. *Cell Death Differ.*, 1999: 6, 644-651.
- [96] Zhivotovsky, B., and Kroemer, G. Apoptosis and genomic instability. *Nature Rev. Mol. Cell Biol.* 2004: 5, 752-762.
- [97] Sorokina N.I. et al. Molecular mechanisms of radiation death of lymphoid cells. Protein poly-ADP-ribosylation in thymocyte fractions. *Radiobiologia*. ;23 (6):739-42.
- [98] Filippovich I., Sorokina N. , Soldatenkov V., Alfyerova T., Trebenok Z. Effect of the inducers of cellular differentiation and of ionizing radiation on thymus lymphocytes: chromatin degradation and programmed cell death. *Int J Radiat Biol Relat Stud Phys Chem Med.* 1988 Apr. 53 (4):617-28.
- [99] Bebeshko V et al. Biological markers of ionizing radiation. *Atomic Medicine.* N1 , (39) – I/II, 2004. pp.85 -104.
- [100] Popov D., Kirchin V., Maliev V., Zlobin V. All Union conference of agriculture radiobiology (USSR). Obninsk, 1990. P. 136-137.
- [101] Popov D., Kirchin V., Saphonova V, Usupova S. Republican Scientific Conference “Theoretical and practical questions of veterinary medicine” (USSR). Kazan, 1990. P. 34.
- [102] Gel Filtration, Principles and Methods Handbook, USA. 181-0221-8, Amersham, Biosciences.
- [103] Maliev V., Kirchin V., Popov D. Antiradiation vaccine. *Herald of VSC of Russian Academy of Science.* 2003. № 3. P. 12-28.
- [104] Maliev V., Popov D., Casey R, Jones J. Mechanisms of Action for an Anti-Radiation Vaccine in reducing the biological impact of high dose and dose-rate, low-linear energy transfer Radiation Exposure. *J. Radiation Biology*, v.47.N3. pp. 286-291.
- [105] Jaiswal M, Zech W, Goos M., Leutbecher C, Ferri A, Zippelius A, Carri M, Nau R, Keller BU (2009). "Impairment of mitochondrial calcium handling in a mtSOD1 cell culture model of motoneuron disease". *BMC Neuroscience* 10: 64.



- [106] Falkow S. Molecular Koch's postulates applied to bacterial pathogenicity--a personal recollection 15 years later. *Nat Rev Microbiol.* 2004 Jan; 2(1):67-72.
- [107] Jones J, Karouia F, Casey R. "Ionizing Radiation as a Carcinogen". In: McQueen C, ed. *Comprehensive Toxicology, Volume 14: Carcinogenesis.* Oxford EN: Elsevier 2010
- [108] Epperly, et al Modulation of Total Body Irradiation Induced Life Shortening by Systemic Intravenous MnSOD-Plasmid Liposome Gene Therapy. *Radiation Research*, 2008 170(4):437-444
- [109] Epperly, M.W., Wang, H., Jones, J., Dixon, T., Montesinos, C.A., and Greenberger, J.S. Antioxidant-Chemoprevention Diet Ameliorates Late Effects of Total Body Irradiation and Supplements Radioprotection by MnSOD-Plasmid Liposome Administration. *Radiat. Res.* 2011 June 175(6):759-765
- [110] Maliev, V, Popov,D, Jones, JA, Casey, RC. Mechanism of action for anti-radiation vaccine in reducing the biological impact of high-dose irradiation *Advances in Space Research* 2007 40:586-590
- [111] Baum S. Erythrocyte stem cell kinetics in postradiation rat. *Armed Forces Radiobiology Research Institute, Defense Atomic Support Agency, Bethesda, Maryland.* May. 1966.
- [112] Mihai R., Doina G. Membrane structural and functional modifications due to oxidative stress induction by irradiation and chemical agents. *European Commission Center of Excellence, InterDisciplinary research and Applications based on Nuclear and Atomic Physics.* Report WP5 ID RAHAP 25-02/2002. pp. 1-8.
- [113] Benderitter M. et al. The cell membrane as a biosensor of oxidative stress induced by radiation exposure: a multiparameter investigation. *Radiation Research: vol.159, No 4*, pp.471-483.
- [114] Dixon D., Darveau R. Lipopolysaccharide Heterogeneity: Innate Host Responses to Bacterial Modification of Lipid A Structure. *J. DenRes* 84(7): 584-595, 2005.
- [115] Moroni M, Coolbaugh T., Lombardini E, Mitchell J, Moccia K, Shelton L, Nagy V, Whitnall M. Hematopoietic radiation syndrome in the Gottingen minipig. *Radiat Res.* 2011 Jul;176(1): 89-101. Epub 2011 Apr 26.
- [116] Kudriashov I., Goncharenko E, Deev L, Gorskaia T, Samořlikova T. Depression of endogenous radiation-sensitizing agents--lipid toxic substances--as one of the mechanisms of radiation protection. *Dokl Akad Nauk SSSR.* 1970; 195(1):206-8.
- [117] Smirnov D., Brady L., Halasa K, Morley M, Solomon S. and Cheun V.. Genetic variation in radiation induced cell death. *Genome Research.* August 15. 2011.
- [118] Sanchez M., Benitez A., Ortloff L., Green L. Alterations in Glutamate Uptake in NT2-Derived Neurons and Astrocytes after Exposure to Gamma Radiation. *Radiation Research.* 2009 January; 171(1): 41-52.



## **Current Topics in Ionizing Radiation Research**

Edited by Dr. Mitsuru Neno

ISBN 978-953-51-0196-3

Hard cover, 840 pages

**Publisher** InTech

**Published online** 12, February, 2012

**Published in print edition** February, 2012

Since the discovery of X rays by Roentgen in 1895, the ionizing radiation has been extensively utilized in a variety of medical and industrial applications. However people have shortly recognized its harmful aspects through inadvertent uses. Subsequently people experienced nuclear power plant accidents in Chernobyl and Fukushima, which taught us that the risk of ionizing radiation is closely and seriously involved in the modern society. In this circumstance, it becomes increasingly important that more scientists, engineers and students get familiar with ionizing radiation research regardless of the research field they are working. Based on this idea, the book "Current Topics in Ionizing Radiation Research" was designed to overview the recent achievements in ionizing radiation research including biological effects, medical uses and principles of radiation measurement.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Dmitri Popov, Jeffrey Jones and Vacheslav Maliev (2012). Radiation Toxins - Effects of Radiation Toxicity, Molecular Mechanisms of Action, Radiomimetic Properties and Possible Countermeasures for Radiation Injury, Current Topics in Ionizing Radiation Research, Dr. Mitsuru Neno (Ed.), ISBN: 978-953-51-0196-3, InTech, Available from: <http://www.intechopen.com/books/current-topics-in-ionizing-radiation-research/radiation-toxins-molecular-mechanisms-of-toxicity-and-radiomimetic-properties->

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen